



31. Let \vec{a}, \vec{b} and \vec{c} be three non-zero vectors such that no two of them are collinear and $(\vec{a} \times \vec{b}) \times \vec{c} = \frac{1}{2} |\vec{b}| |\vec{c}| |\vec{a}|$. If θ is the angle between vectors \vec{b} and \vec{c} , then a value of $\sin \theta$ is

1. $\frac{2}{3}$
2. $\frac{-2\sqrt{3}}{3}$
3. $\frac{2\sqrt{2}}{3}$
4. $\frac{-\sqrt{2}}{3}$

Solution: (3)

$$(\vec{a} \cdot \vec{c}) \vec{b} - (\vec{b} \cdot \vec{c}) \vec{a} = \frac{1}{3} (|\vec{b}| |\vec{c}|) \vec{a}$$

comparing $\cos \theta = -\frac{1}{3} \Rightarrow \sin \theta = \frac{2\sqrt{2}}{3}$


32. Let O be the vertex and Q be any point on parabola, $x^2=8y$. if the point P divides the line segment OQ internally in the ratio 1:3, then the locus of P is :

1. $y^2 = 2x$
2. $x^2 = 2y$
3. $x^2 = y$
4. $y^2 = x$

Solution: (2)

$$x = \frac{4t}{4} = t$$

$$y = \frac{2t^2}{4} = \frac{t^2}{2}$$



so, $x^2 = 2y$ is locus of p.


33. If the angles of elevation of the top of a tower from three collinear points, A, B, and C, on a line leading to the foot of the tower, are 30° , 45° and 60° respectively, then the ratio, AB : BC, is :

1. $1:\sqrt{3}$
2. $2:3$
3. $\sqrt{3}:1$
4. $\sqrt{3}:\sqrt{2}$

Solution: (3)

$$\frac{AQ}{\sqrt{3}} = BQ = CQ\sqrt{3}$$

$$\therefore AQ = 3CQ, AC = 2CQ$$



$$AB = (3 - \sqrt{3})CQ = \sqrt{3}(\sqrt{3} - 1)CQ$$

$$\therefore AB : BC = \frac{\sqrt{3}(\sqrt{3} - 1)CQ}{(\sqrt{3} - 1)CQ} = \frac{\sqrt{3}}{1}$$

34. The number of points, having both co-ordinates as integers, that lie in the interior of the triangle with vertices (0,0), (0, 41) and (41,0), is:


1. 820
2. 780
3. 901
4. 861

Solution: (2)

number of points

$$= 1+2+3+ \dots + 39$$

$$= \frac{39 \times 40}{2} = 780$$



35. the equation of the plane containing the line $2x - 5y + z = 3$; $x + y + 4z = 5$, and parallel to the plane, $x + 3y + 6z = 1$ is :

- $x + 3y + 6z = 7$
- $2x + 6y + 12z = -13$
- $2x + 6y + 12z = 13$
- $x + 3y + 6z = -7$

Solution: (1)

equation of required plane is

$$2x - 5y + z - 3 + \lambda(x + y + 4z - 5) = 0$$

$$\Rightarrow (2 + \lambda)x + (\lambda - 5)y + (1 + 4\lambda)z - 3 - 5\lambda = 0 \dots\dots(1)$$

Since, Required plane is parallel to

$$x + 3y + 6z = 1,$$

$$\text{So, } \frac{2 + \lambda}{1} = \frac{\lambda - 5}{3} = \frac{1 + 4\lambda}{6} = \frac{3 + 5\lambda}{1}.$$

Solving any two pair, we get $\lambda = 11/2$.

Putting value of λ in equation (1), we get

$$x + 3y + 6z = 7$$

36. Let A and B be two sets containing four and two elements respectively. Then the number of subsets of the set $A \times B$, each having at least three elements is:

- 275
- 510
- 219
- 256

Solution: (3)

number of subsets =


$$2^8 - \{c(8,0) + c(8,1) + c(8,2)\} = 256 - (1 + 8 + 28) = 219.$$

37. Locus of the image of the point (2,3) in the line $(2x - 3y + 4) + k(x - 2y + 3) = 0$,

$k \in \mathbb{R}$, is a:

- circle of radius $\sqrt{2}$.
- circle of radius $\sqrt{3}$.
- straight line parallel to x-axis.
- straight line parallel to y-axis.

Solution: (1)



Clearly, the locus of Q is a circle.

38. $\lim_{x \rightarrow 0} \frac{(1 - \cos 2x)(3 + \cos x)}{x \tan 4x}$ is equal to:

- 2
- $\frac{1}{2}$
- 4
- 3

Solution: (1)

$$\lim_{x \rightarrow 0} \frac{(1 - \cos 2x)}{x^2} \times \frac{(3 + \cos x)}{\frac{\tan 4x}{x}}$$

$$= \frac{4}{2} \times \frac{4}{2} = 2$$

39. The distance of the point (1,0,2) from the point of intersection of the line $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12}$ and the plane $x - y + z = 16$ is:

- $3\sqrt{21}$
- 13
- $2\sqrt{14}$
- 8

Solution: (2)

Equation of line is $\frac{x-2}{3} = \frac{y+1}{4} = \frac{z-2}{12} = \lambda$ (say)

$$\Rightarrow x = 3\lambda + 2, y = 4\lambda - 1, z = 12\lambda + 2$$

Putting these values in plane $x - y + z = 16$, we get, $\lambda = 1$.

So, point of intersection (5,3,14)

distance = $\sqrt{4^2 + 3^2 + 12^2} = 13$

40. The sum of coefficients of integral powers of x in the binomial expansion of $(1 - 2\sqrt{x})^{50}$ is

- $\frac{1}{2}(3^{50} - 1)$
- $\frac{1}{2}(2^{50} + 1)$
- $\frac{1}{2}(3^{50} + 1)$
- $\frac{1}{2}(3^{50})$

Solution: (3)

Let $\sqrt{x} = y$.

Now,

$$(1 - 2y)^{50} = a_0 + a_1y + a_2y^2 + \dots + a_{50}y^{50} \dots\dots(1)$$

Putting $y = 1$, in (1) we get

$$1 = a_0 + a_1 + a_2 + \dots + a_{50} \dots\dots(2)$$

putting $y = -1$, in (1) we get,

$$3^{50} = a_0 - a_1 + a_2 - a_3 \dots + a_{50} \dots (3)$$

Adding (2) and (3), we get

$$3^{50} + 1 = 2(a_0 + a_2 + \dots + a_{50})$$

$$\Rightarrow (a_0 + a_2 + \dots + a_{50}) = \frac{3^{50} + 1}{2}$$

41. the sum of first 9 terms of the series

$$\frac{1^3}{1} + \frac{1^3 + 2^3}{1+3} + \frac{1^3 + 2^3 + 3^3}{1+3+5} + \dots \text{ is:}$$

1. 14
2. 192
3. 71
4. 96

Solution: (4)

$$T_n = \frac{(n+1)^2}{4} = \frac{(n^2 + 2n + 1)}{4}$$

$$s_n = \sum T_n = \frac{1}{9} \left[\frac{n(n+1)(2n+1)}{6} + 2 \frac{(n)(n+1)}{2} + n \right]$$

$$= \frac{n}{24} (2n^2 + 9n + 13)$$


$$\therefore s_9 = \frac{9}{24} \times (2 \times 81 + 81 + 13) = 96$$

42. The area (in sq. units) of the region described by

$$\{(x, y) : y^2 \leq 2x \text{ and } y \geq 4x-1\}$$
 is:

1. $\frac{15}{64}$
2. $\frac{9}{32}$
3. $\frac{7}{32}$
4. $\frac{5}{64}$

Solution: (2)



$$\text{Area} = \int_{y=-1/2}^{y=1} \left(\frac{y+1}{4} - \frac{y^2}{2} \right) dy = 9/32$$

43. The set of all values of λ for which the system of linear equations:

$$2x_1 - 2x_2 + x_3 = \lambda x_1$$

$$2x_1 - 3x_2 + 2x_3 = \lambda x_1$$

$$-x_1 + 2x_3 = \lambda x_3$$

has a non-trivial solution,

1. contains two elements
2. contains more than two elements.
3. is an empty set
4. is singleton.

Solution: (1)

$$(2 - \lambda) x_1 - 2x_2 + x_3 = 0$$

$$2x_1 - (3 + \lambda) x_2 + 2x_3 = 0$$

$$-x_1 + 2x_2 - \lambda x_3 = 0$$

$$\begin{vmatrix} 2-\lambda & -2 & 1 \\ 2 & -(3+\lambda) & 2 \\ -1 & 2 & -\lambda \end{vmatrix} = 0$$

$$\Rightarrow (\lambda - 1)^2 (\lambda + 3) = 0$$

$$\Rightarrow \lambda = 1 \text{ and } \lambda = -3.$$

So, set contains two elements.

44. a complex number z is said to be unimodular if $|z|=1$. Suppose z_1 and z_2 are complex number

such that $\frac{z_1 - 2z_2}{2 - z_1 z_2}$ is unimodular and z_2 is not unimodular. Then the point z_1 lies on a :

1. circle of radius 2.
2. circle of radius $\sqrt{2}$.
3. straight line parallel to x-axis.
4. straight line parallel to y-axis.

Solution: (1)

$$\left| \frac{z_1 - 2z_2}{2 - z_1 z_2} \right| = 1 \Rightarrow |z_1 - 2z_2|^2 = |2 - z_1 z_2|^2$$

$$\Rightarrow (z_1 - 2z_2)(\overline{z_1 - 2z_2}) = (2 - z_1 z_2)(\overline{2 - z_1 z_2})$$

$$\Rightarrow (|z_1|^2 - 4)(1 - |z_2|^2) = 0$$

$$\Rightarrow |z_1|^2 = 4, \therefore z_1 \text{ is not unimodular}$$


$$\Rightarrow |z_1| = 2 \text{ which lies on a circle of radius 2.}$$

45. The number of common tangents to the circles
 $x^2 + y^2 - 4x - 6y - 12 = 0$

and $x^2 + y^2 + 6x + 18y + 26 = 0$ is :

1. 3 2. 4
 3. 1 4. 2

Solution : (1)



$$x^2 + y^2 - 4x - 6y - 12 = 0$$

$$C_1(2,3), r_1 = \sqrt{4 + 9 + 12} = 5$$

$$x^2 + y^2 + 6x + 18y + 26 = 0$$

$$C_2(-3,-9), r_2 = \sqrt{9 + 81 - 26} = 8$$

$$\overline{C_1C_2} = \sqrt{25 + 144} = 13$$

46. The number of integers greater than 6,000 that can be formed, using the digits, 3,5,6, 7 and 8, without repetition, is:

1. 120 2. 72
 3. 216 4. 192

Solution: (4)

$$\text{Total number of integers} = 3(24) + 5!$$

$$= 72 + 120 = 192.$$

47. Let $y(x)$ be the solution of the differential equation

$$(x \log x) \frac{dy}{dx} + y = 2x \log x, (x \geq 1).$$

Then $y(e)$ is equal to:

1. 2 2. $2e$
 3. e 4. 0

Solution:

The question is not theoretically correct.

48. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -1 \\ a & 2 & b \end{bmatrix}$ is a matrix satisfying the

equation $AA^T = 9I$, where I is 3×3 identity matrix, then the ordered pair (a,b) is equal to:

1. $(2,1)$ 2. $(-2, -1)$
 3. $(2,-1)$ 4. $(-2,1)$

Solution:(2)

$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ 9 & 2 & 6 \end{bmatrix} \begin{bmatrix} 1 & 2 & 9 \\ 2 & 1 & 2 \\ 2 & -2 & 6 \end{bmatrix} = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 9 \end{bmatrix}$$

Equating the corresponding elements, we get,

$$a + 2b = -4 \text{ and } a - b = -1.$$

Solving both equations, we get

$$a = -2 \text{ and } b = -1.$$

49. If m is the A.M. of two distinct real numbers l and n ($l, n > 1$) and G_1, G_2 and G_3 are three geometric means between l and n , then

$G_1^4 + 2G_2^4 + G_3^4$ equals.

1. $4lmn^2$ 2. $4l^2m^2n^2$
 3. $4l^2mn$ 4. $4lm^2n$

Solution: (4)

$$l + n = 2m.$$

l, G_1, G_2, G_3, n are in G.P.

$$\Rightarrow \text{common ratio} = r = \left(\frac{n}{l}\right)^{1/4}$$

$$\Rightarrow G_1^4 = l^3n, G_2^4 = l^2n^2, G_3^4 = ln^3$$

$$\Rightarrow G_1^4 + 2G_2^4 + G_3^4 = l^3n + 2l^2n^2 + ln^3$$

$$= \ln(l+n)^2 = \ln(4m^2) = 4lm^2n.$$

50. the negation of $\sim s \wedge (\sim r \vee s)$ is equivalent to:

1. $s \wedge (r \vee \sim s)$ 2. $s \wedge r$
 3. $s \wedge \sim r$ 4. $s \wedge (r \wedge \sim s)$

Solution:(2)

$$\sim[\sim s \vee (\sim r \wedge s)] = \sim((\sim s \vee \sim r) \wedge (\sim s \vee s))$$

$$= \sim(\sim s \vee \sim r) = (s \wedge r).$$

51. The integral $\int \frac{dx}{x^2(x^4 + 1)^{3/4}}$ equals:

1. $-(x^4 + 1)^{\frac{1}{4}} + c$ 2. $-\left(\frac{x^4 + 1}{x^4}\right)^{\frac{1}{4}} + c$
3. $\left(\frac{x^4 + 1}{x^4}\right)^{\frac{1}{4}} + c$ 4. $(x^4 + 1)^{\frac{1}{4}} + c$

Solution:(2)

$$\int \frac{dx}{x^2(x^4 + 1)^{3/4}} = \int \frac{dx}{x^5(1 + \frac{1}{x^4})^{3/4}}$$

$$= \int \frac{(-t^3)dt}{(t^3)^3} \left\{ \text{put } 1 + \frac{1}{x^4} = t^4 \Rightarrow \frac{1}{x^5} dx = -t^3 dt \right.$$

$$= -t + C = -\left(\frac{x^4 + 1}{x^4}\right)^{1/4} + C$$

52. The normal to the curve, $x^2 + 2xy - 3y^2 = 0$, at (1,1):

- meets the curve again in the third quadrant.
- meets the curve again in the fourth quadrant.
- does not meet the curve again.
- meets the curve again in the second quadrant.

Solution:(2)

$$x^2 + 2xy - 3y^2 = 0 \Rightarrow (x - y)(x + 3y) = 0$$

$$\Rightarrow x = y \text{ or } x + 3y = 0.$$

(1,1) lies on the line $x = y$.

Equation of normal to it is $y - 1 = (-1)(x - 1)$

$$\Rightarrow x + y = 2.$$

This normal meets the curve $x + 3y = 0$ at (3,-1) which lies in 4th quadrant.

53. Let

$$\tan^{-1} y = \tan^{-1} x + \tan^{-1} \left(\frac{2x}{1 - x^2} \right)$$

where $|x| < \frac{1}{\sqrt{3}}$. Then a value to y is:

- $\frac{3x - x^3}{1 + 3x^2}$
- $\frac{3x + x^3}{1 + 3x^2}$
- $\frac{3x - x^3}{1 - 3x^2}$
- $\frac{3x + x^3}{1 - 3x^2}$

Solution : (3)

$$\tan^{-1} y = \tan^{-1} x + \tan^{-1} \left(\frac{2x}{1 - x^2} \right)$$

$$\Rightarrow \tan^{-1} y = \tan^{-1} x + 2 \tan^{-1} x$$

$$\Rightarrow \tan^{-1} y = 3 \tan^{-1} x$$

$$\Rightarrow \tan^{-1} y = \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right)$$

$$\Rightarrow y = \left(\frac{3x - x^3}{1 - 3x^2} \right)$$

54. If the function.

$$\begin{cases} k\sqrt{x+1}, & 0 \leq x \leq 3 \\ mx + 2, & 3 < x \leq 5 \end{cases}$$

is differentiable, then the value of k+m is:

- $\frac{10}{3}$
- 4
- 2
- $\frac{16}{5}$

Solution: (3)

$$g(x) = \begin{cases} k\sqrt{x+1}, & 0 \leq x \leq 3 \\ mx + 2, & 3 < x \leq 5 \end{cases}$$

$g(x)$ is differentiable

$$\Rightarrow g'(3^-) = g'(3^+) \Rightarrow \left(\frac{k}{2\sqrt{x+1}} \right)_{x=3} = m$$

$$\Rightarrow k = 4m \dots\dots\dots(1)$$

Next, $g(x)$ is continuous as it is differentiable.

$$\text{So, } g(3^-) = g(3^+) \Rightarrow 2k = 3m + 2 \dots\dots\dots(2)$$

From (1) and (2),

$$\text{we get } m = 2/5 \Rightarrow k + m = 2$$

55. The mean of the data set comprising of 16 observations is 16. If one of the observations valued 16 is deleted and three new observations valued 3, 4 and 5 are added to the data, then the mean of the resultant data, is

1. 15.8
2. 14.0
3. 16.8
4. 16.0

Solution: (2)

$$\text{Sum} = S = 16 \times 16 = 256.$$

$$\text{Mean} = \frac{S - 16 + 3 + 4 + 5}{18} = \frac{256 - 16 + 12}{18} = 14.$$

56. The integral

$$\int_2^4 \frac{\log x^2}{\log x^2 + \log(36 - 12x + x^2)} dx \text{ is equal to:}$$

1. 1
2. 6
3. 2
4. 4

Solution: (1)

$$I = \int_2^4 \frac{\log x^2}{\log x^2 + \log(36 - 12x + x^2)} dx$$

$$= \int_2^4 \frac{\log x^2}{\log x^2 + \log(6-x)^2} dx$$

$$= \int_2^4 \frac{\log(6-x)^2}{\log(6-x)^2 + \log x^2} dx$$

$$\Rightarrow 2I = \left[X \right]_2^4 \Rightarrow I = 1.$$

57. Let α and β be the roots of equation

$$x^2 - 6x - 2 = 0. \text{ If } a_n = \alpha^n - \beta^n, \text{ for } n \geq 1, \text{ then}$$

the value of $\frac{a_{10} - 2a_8}{2a_9}$ is equal to :

1. 3
2. -3
3. 6
4. -6

Solution: (1)

$$x^2 - 6x - 2 = 0 \Rightarrow \alpha + \beta = 6, \alpha\beta = -2$$

$$\text{Now, } \frac{a_{10} - 2a_8}{2a_9} = \frac{(\alpha^{10} - \beta^{10}) - 2(\alpha^8 - \beta^8)}{2(\alpha^9 - \beta^9)}$$

$$\Rightarrow \frac{(\alpha^{10} - \beta^{10}) + \alpha\beta(\alpha^8 - \beta^8)}{2(\alpha^9 - \beta^9)} = \frac{\alpha^9(\alpha + \beta) - \beta^9(\alpha + \beta)}{2(\alpha^9 - \beta^9)}$$

$$= \frac{(\alpha^9 - \beta^9)(\alpha + \beta)}{2(\alpha^9 - \beta^9)} = \frac{(\alpha + \beta)}{2} = 3.$$

58. Let $f(x)$ be polynomial of degree four having extreme values at $x=1$ and $x=2$.

If $\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{x^2} \right] = 3$, then $f(2)$ is equal to :

1. 0
2. 4
3. -8
4. -4

Solution: (1)

$$\text{Let } f(x) = x^2(ax^2 + bx + c).$$

$$\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{x^2} \right] = 3 \Rightarrow c + 1 = 3 \Rightarrow c = 2.$$

$$\text{Again, } f(x) = x^2(ax^2 + bx + c)$$

$$\Rightarrow f'(x) = x^2(2ax + b) + 2x(ax^2 + bx + c)$$

$$\Rightarrow f'(x) = 4ax^3 + 3bx^2 + 2cx$$

$$\text{We have, } f'(1) = 0 \Rightarrow 4a + 3b + 4 = 0 \dots\dots(1)$$

$$\text{and } f'(2) = 0 \Rightarrow 32a + 12b + 8 = 0 \dots\dots(2)$$

From (1) and (2) we have,

$$a = 1/2 \text{ and } b = -2$$


$$\therefore f(2) = 0.$$

59. The area (in sq. units) of the quadrilateral formed by the tangents at the end points of the latera recta to the ellipse

$$\frac{x^2}{9} + \frac{y^2}{5} = 1 \text{ is}$$

1. $\frac{27}{2}$
2. 27
3. $\frac{27}{4}$
4. 18

Solution: (2)



Equation of tangent at $\left(ae, \frac{b^2}{a} \right)$ is given by,

$$\frac{aex}{a^2} + \frac{b^2y}{ab^2} = 1 \Rightarrow \frac{ex}{a} + \frac{y}{a} = 1$$

The tangent meets the axes at $A(a/e, 0)$ and $B(0, a)$.

Similarly we can find out $C(-a/e, 0)$ and $D(0, -a)$.

$$\text{Area of quadrilateral} = \frac{1}{2} \times 2a \times \frac{2a}{e} = \frac{2a^2}{e}$$

$$= \frac{(2)(9)}{(2/3)} = 27 \quad \left\{ \begin{array}{l} \because 9(1 - e^2) = 5 \\ \Rightarrow e^2 = 1 - \frac{5}{9} = \frac{4}{9} \Rightarrow e = 2/3 \end{array} \right.$$

60. If 12 identical balls are to be placed in 3 identical boxes, then the probability that one of the boxes contains exactly 3 balls is:

1. $220 \left(\frac{1}{3} \right)^{12}$ 2. $22 \left(\frac{1}{3} \right)^{11}$

3. $\frac{55}{3} \left(\frac{2}{3} \right)^{11}$ 4. $55 \left(\frac{2}{3} \right)^{10}$

Solution: Theoretically the question is wrong.

